

Efficacy of Lycopene Capsules in Oral Submucous Fibrosis

¹Dr. Samar Ali Faraz, ²Dr. Pratik Parkarwar, ³Dr. Prashant Kore, ⁴Dr. Shruti Wadne ,

⁵Dr. Vasundhara Bandagi, ⁶Dr. Usama Kharadi

^{1,3}MDS, ^{2,6}Asso. Professor, ^{4,5}P.G Student

^{1, 2, 4, 5, 6} Dept of Oral Medicine & Radiology, & Radiology PDU Dental College, Solapur.

³Dept. of Periodontology & Implantology

Corresponding Author

Dr. Pratik Parkarwar

Asso. Professor. Dept of Oral Medicine & Radiology. PDU Dental College, Solapur.

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

ABSTRACT

Oral sub mucous fibrosis is the premalignant condition affecting the oral cavity and most prevailing in youthful generation. The etiology of this condition is consumption of betel nut causing reduce mouth opening, burning sensation and finally the fibrosis of mucosa. The subjects have the symptoms like stiffness of oral mucosa, difficulty in swallowing, alter speech, shrunken uvula. So this condition can be prevented and controlled by stoppage of computation of betel nut , topical and systemic medicines , mouth opening exercises and surgery .In the medicines antioxidant, corticosteroids (topical),turmeric, immused milk, vasodilators are used and found to be effective. The study was undertaken on 50 subjects suffering from Grade I and Grade II OSMF with fifteen days interval and prescribed on Lycopene and placebo for three months. There was significant improvement in burning sensation and mouth opening of OSMF subjects.

The study concluded that the administration of Lycopene capsules showed improvement in mouth opening and decrease in the burning sensation in the oral cavity.

Keywords: Oral Submuocus Fibrosis, Premalignant condition, Lycopene Capsules, Stiffness, Burning, Trismus, Betel nut.

INTRODUCTION

Oral Submucous Fibrosis is defined as an insidious, chronic disease which affects any part of the oral cavity and sometimes the pharynx¹ and is occasionally preceded by and/or associated with vesicle formation² and is always associated with a juxta-epithelial inflammatory reaction which is followed by progressive hyalinization of the lamina propria³ leading to stiffness of the oral mucosa and deeper tissues with progressive limitation in opening of the mouth and protrusion of the tongue leading to difficulty in eating, swallowing and phonation.^{3,4}

It is a precancerous condition seen most commonly in the Indian sub-continent and has a reported incidence of between 0.2–1.2% of the urban population who attend the dental clinic and the condition shows a

female to :male predilection of 3:1 and characteristically first presents in adulthood between the ages of 45–54 years.⁵The etiology is multifactorial origin for this condition and the various hypothesis implicated include the role of local irritants such as capsaicin⁶, tobacco⁷, areca nut⁸⁻¹², pungent and spicy foods¹³, and alcohol¹⁴, iron and vitamin B-complex deficiency, anaemia¹⁵, and a genetic predisposition to the disease¹².

There are different treatment modalities for this condition, here one such treatment modality in the form of administration of antioxidants in the form of Lycopene capsules was carried out on 50 patients who attended the Outpatient Department of OMR.

Aim of the study: To assess the efficacy of Lycopene capsules in the management of patients with Oral Submucous Fibrosis.

Source of data.

Fifty (50) adult patients who enrolled with signs and symptoms of OSMF were included in the study. Following parameters were included in the establishment of diagnosis and these 2 parameters were satisfied for inclusion in the study. A. Positive history of chewing of areca nut or one of its commercial preparations, difficulty in chewing and swallowing, and having burning sensation on eating spicy food. B. Restricted mouth opening

Following establishment of diagnosis, each patient was informed about the condition, its precancerous potential and advised to discontinue use of areca nut in all forms. A detailed case history including habit of history with details of duration, in years, frequency of chews per day was taken. All patients underwent oral prophylaxis to remove extrinsic stains, in order to motivate the patient towards recovery and to inform the investigator if patient resumes habit. Each patient was screened for diagnosis and inclusion, examined on three occasions, day 1 which is the first day of starting treatment, day 30, day 60, day 90 which is the fourth and last day of evaluation. Each

patient was administered Lycopene 4 mg capsules, 2 capsules twice a day for 90 days.

Method of data collection

Patients were evaluated for the following criteria


1. Mouth opening based on interincisal separation: Distance between the upper and lower central incisors when maximally extended with mouth wide open. Normal values: Males-35-45 mm. Females-30-42 mm.



Figure

2. Burning sensation present or not and if so, degree to be determined by use of a Verbal Analogue Scale.(VAS)

VERBAL ANALOGUE SCALE FOR BURNING SENSATION -



8 - Continuous

9 - Severe

10 - Intolerably severe

Inclusion Criteria:

Patients suffering from chronic oral mucous fibrosis lesions characterized by burning sensations in the mouth, particularly while taking hot and spicy foods. Adult patients of either sex aged between 18-50 years were included. Patients who had not participated in a similar investigation in past four weeks. Patient willing to give a written informed consent and follow the schedule.

Exclusion criteria:

Patients suffering from severe systemic disorders. Age below 18 years. Patients who refused to sign informed consent. Patients having a known history or present condition of allergic response to similar pharmaceutical products, pre-existing systemic disease necessitating long-term medications and pregnant and lactating women were excluded from the study.

Method of data analysis.

Results were analyzed statistically by means of a paired t-test analysis.

Results:

TABLE NO.1: DISTRIBUTION OF MEAN AND SD VALUES OF MOUTH OPENING (MM) AT DAY 0, AT DAY 30, AT DAY 60 AND AT DAY 90:

Duration	Mouth Opening(mm) (n=30)	Student's Paired 't' test value	'p' value	Significance
	Mean \pm SD			
Day 0	28.19 \pm 7.80	50.11	p<0.01	Highly significant
Day 30	29.23 \pm 7.86			
Day 60	30.04 \pm 8.02			
Day 90	30.66 \pm 8.07			

By applying Student's Paired 't' test there is a highly significant increase in mean mouth opening (mm) from starting treatment i.e. day 0 to last day of evaluation i.e. day 90 (p<0.01). The average increase is 2.46 mm \pm 1.08 mm.

TABLE NO.2: DISTRIBUTION OF MEAN AND SD VALUES OF BURNING SENSATION (VAS) AT DAY 0, AT DAY 30, AT DAY 60 AND AT DAY 90:

Duration	Burning Sensation (VAS) (n=30)	Student's Paired 't' test value	'p' value	Significance
	Mean \pm SD			
Day 0	2.97 \pm 0.96	73.26	p<0.01	Highly significant
Day 30	2.77 \pm 0.93			
Day 60	2.27 \pm 0.90			
Day 90	2.03 \pm 0.88			

By applying Student's Paired 't' test there is a highly significant decrease in mean **Burning Sensation (VAS)** from starting treatment i.e. day 0 to last day of evaluation i.e. day 90 ($p < 0.01$). The average decrease is $0.94 \text{ mm} \pm 0.02 \text{ mm}$.

Discussion:

Oral submucous fibrosis is a precancerous condition affecting the buccal mucosa of the oral cavity most commonly leading to marked rigidity and inability to eat^{16,17} however any part of the oral cavity may be involved including the pharynx¹⁸ eventually leading to decrease in mouth opening. Malignant transformation rate of OSMF is as high as 7.6% in a study conducted in the Indian subcontinent over a 17 year period.¹⁹ Treatment options include initially stoppage of habit followed by avoidance of spicy food and ingestion of chillies followed by intralesional steroid injections, placental extracts, pentoxifylline, lycopene, surgical excision and laser therapy¹⁷ can be used to a greater effect in reducing the signs and symptoms of OSMF.

Lycopene is an effective antioxidant from tomato extract and has been proved to be the most potent radical scavenger. It has the mechanism of action like antioxidant activity, inhibition of cancer cell proliferation, restoration of gap junctions, regulation of transcription, interfere with growth factor. It was successfully tried in leukoplakia where it has given excellent and favorable results. It is also used in lichen planus, oral cancer, gingivitis, diabetes, AIDS and periodontitis. It is a potent anticarcinogenic and has demonstrated profound benefits in precancerous lesions and conditions. Abhinav Kumar et al that there was marked improvement in mouth opening and in noticeable reduction of burning sensation in OSMF patients with help of lycopene treatment than patients treated with placebo. This curative effect of lycopene may be due to inhibition of abnormal fibroblasts, up regulation of lymphocyte resistance to stress and suppression of inflammatory response.¹⁶

Sunderaj Et al (2012) conducted a study in subjects receiving 16 mg lycopene in divided doses and assessed bimonthly for four months and found overall average improvement in mouth opening was 2 mm. There was reduced burning sensation. As compared to our study we done study on 50 subjects receiving 16 mg lycopene in divided doses for three

months with bimonthly follow up. In our study we found that there was reduced burning sensation in all subjects and improvement in mouth opening.¹⁷

Summary and Conclusion:

In this study the efficacy of Lycopene capsules was evaluated in 50 OSMF patients. Lycopene capsules due to its special antioxidant properties were found to be effective in the improvement of signs and symptoms of Oral Submucous Fibrosis. It is effective in reducing the objective signs of OSMF thereby improving the mouth opening (percentage of which was 8.05%), and decreasing the burning sensation (percentage of which was 31.54%). Thus, it can be concluded that Lycopene capsules appear to be a very promising drug in the improvement of the symptoms and management of Oral Submucous Fibrosis.

References:

1. Paul R R, Mukherjee A, Dutta PK, Banerjee S, Pal M, Chatterjee J.A. Novel wavelet neural network based pathological stage detection technique for an oral precancerous condition. *J Clin Pathol*. 2005; 58(9):932-8.
2. Pindborg JJ, Sirsat SM. Oral submucous fibrosis. *Oral Surg Oral Med Oral Pathol*. 1966; 22: 764-79.
3. Reddy V, Wanjari PV. Oral submucous fibrosis: Correlation of clinical grading to various habit factors. *Int J Dent Clin*. 2011; 3: 21-4.
4. Schwartz J. Atrophia idiopathica (tropica) mucosa oris. Demonstrated at the 11th International Dental Congress, London. 1952.
5. Joshi SG. Submucous fibrosis of the palate and pillars. *Indian J Otolaryngol*. 1953; 4: 14.
6. Lal D. Diffuse oral submucous fibrosis. *J All India Dent Assoc*. 1953; 26: 1-3, 14-15.
7. Su JP. Idiopathic scleroderma of the mouth. Report of three cases. *Arch Otolaryngol*. 1954; 59: 330-2.
8. Rao ABN. Idiopathic palatal fibrosis. *Br J Surg* 1%. 1962; 50: 23-5.
9. Khan S, Chatra L, Prashanth SK, Veena KM, Rao PK. Pathogenesis of oral submucous fibrosis. *J Cancer Res Ther*. 2012; 8: 199-203.

10. Guta MK, Mhaske S. Oral submucous fibrosis: Current concepts in etioathogenesis. *People's J Sci Res*. 2008; 1:39-44.
11. Chang YC, Hu CC, Tseng TH, Tai KW, Lii CK, Chou MY. Synergetic effects of nicotine on arecoline-induced cytotoxicity in human buccal mucosal fibroblasts. *J Oral Pathol Med*. 2001; 30: 458-64.
12. Manoharan S, Kolanjiappan K, Suresh K, Panjamurthy K. Lipid peroxidation and antioxidants status in patients with oral squamous cell carcinoma. *Indian J MedRes*. 2005; 122: 529-34.
13. Areca nut. Taipei Times of oral pathology and medicine 23 (2): 65-9.
14. WHO IARC monograph program finds betelquid and areca nut chewing carcinogenic to human. GENEVA, 2003.
15. Murti PR, Bhonsle RB, Gupta PC, Daftary DK, Pindborg JJ, Metha FS. Aetiology of oral submucous fibrosis with special reference to the role of areca nut chewing. *J Oral Pathol Med*. 1995;24:145-52.
16. Abhinav Kumar, Anjana Bagewadi, Vaishali Keluskar, and Mohitpal Singh, Efficacy of lycopene in the management of oral submucous fibrosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;103:207-13.
17. S Sunder raj, Rohit Sharma, Vishal Agarwal, Praful Narang, Reddy Yg, Amit Kumar Sharma. An invivo study to determine the efficacy of lycopene as compared to multivitamin preparation in the treatment of oral submucous fibrosis. *JIAOMR*. 2012;24(3):190-93