



Knowledge, Attitude, and Behavior of Breastfeeding among Urban Women in Chennai- A Cross-Sectional Study

¹K.Saravanan, ²S.Hariharan, ³Deepika.P, ⁴B.Krishna Prasanth

¹Associate Professor, ^{2,3,4}Assistant Professor, ⁴Epidemiologist

^{1,3}Department of Pediatrics, ^{2,4}Department of Community Medicine,

¹Venkateshwara Institute of Medical Sciences, Gajraula, Uttar Pradesh

^{2,3,4}Sree Balaji Medical College & Hospital, Bharath Institute of Higher Education & Research, Chennai, Tamilnadu

***Corresponding Author:**

Dr. B. Krishna Prasanth

Epidemiologist & Assistant Professor, Department of Community Medicine,
Sree Balaji Medical College & Hospital, Bharath Institute of Higher Education & Research, Chennai

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract

Background

Appropriate breastfeeding practices play a prominent role in reducing morbidity and mortality among children below 5 years of age. Exclusive breastfeeding is important for a child's health and growth. However, there seems to be a disdain for breastfeeding practices among emergent countries. This study intends to assess the knowledge, attitude, and behavior towards breastfeeding among nursing mothers from Chennai.

Methods

This was a cross-sectional study conducted among 200 mothers of children aged 0–1 years. The participants were selected through purposive sampling from postnatal OPD of government hospitals in Chennai. Breastfeeding practices were assessed using a self-administered questionnaire after getting informed consent.

Results

Almost 52% of the mothers had fair knowledge and only a minimal (3%), had poor knowledge regarding breastfeeding practices. Most of the participants 173 (86.5%) believed that adequate feeding with breast milk played an important role in the long-term health of their children. Most of the mothers (77.5%) viewed breastfeeding as more convenient than formula feeding and around 73% of them felt that breastfeeding promoted bonding between the mother and their infants. Almost 72.5% of the study participants initiated breastfeeding as soon as they delivered. Only minimal group of babies (7%) had received pre-lacteal feeds, the commonest being sugar water and honey. Almost 90.5% of new-borns received colostrum as their first feed.

Conclusion

This study inferred that there was only a fair knowledge about breastfeeding among the nursing mothers living in urban regions and also their attitude and behaviour about breastfeeding were sub-optimal. This study also gives an insight into factors hindering optimal breastfeeding practices among young mothers. It is absolutely imperative that we provide proper information, education and communication to nursing mothers regarding proper breast feeding. Henceforth this study identifies the need for public health education campaigns towards breastfeeding practices among nursing mothers living in emergent countries.

Keywords: breastfeeding, women, knowledge

Introduction

Breastfeeding is absolutely indispensable in not only providing the young infant with the necessary nutrients they need for optimal growth and development, but also promotes a direct bond between the mother and her infant(1). Exclusive breastfeeding (EBF) is the choice of nutrition till 6 months, beyond which complementary foods are introduced along with breast feeding. Ideally, breastfeeding is to be initiated within an hour after birth. WHO recommends feeding colostrum, the first milk which is rich in antibodies as an initial food for the neonate. Following EBF for six months, solid foods such as mashed vegetables and fruits should be started to complement breastfeeding till two years. Demand breastfeeding must be advocated and the use of bottles or pacifiers should be discouraged. Breastfeeding has numerous advantages for both the mother and the neonate. Exclusive breastfeeding is related to decreased mortality and morbidity in under 5 children. Breast milk which is easily available and affordable is easily digestible by the developing gastrointestinal system, in comparison with formula feeds. Breast milk meets all the required nutrients necessary during the new-born period, especially in the initial six months.

Under the NHFS survey, among the south Indian states, the state with the least compliance with exclusive breastfeeding was Tamil Nadu in regards to the previous survey. The survey also has come up with results of only 41% of children are breastfed in the first 30 minutes of life, which implies a lost opportunity to give colostrum for the new-born(3). Malnutrition, a more prevalent condition could be eradicated among children if they are breastfed. Children who were breastfed showed higher developmental scores as opposed to their peer groups who were formula-fed. During episodes of diarrhoea, dehydration could be prevented if the child is breastfed. The secretory antibodies present in breast milk reduced the incidence of childhood illnesses such as diarrhoea and pneumonia, the most important causes of infant mortality(5). Maternal benefits from breastfeeding are natural (though not fail-safe) methods of birth control. It reduced the incidence of breast and ovarian cancer. Beyond the immediate benefits, mothers who breastfed were less probably to be obese and its associations. With the promotion of timely breastfeeding, neonatal death accounting for 16% among the child survival

millennium development goal could be achieved if all children were breastfed since the first day and 22% if breastfeeding commenced within the initial 60 minutes(4). Hence this study was undertaken to assess the Knowledge, attitude, and practices of breastfeeding among urban mothers from Chennai.

Materials And Methods

A descriptive cross-sectional study was conducted to assess the knowledge, attitude, and practices of breastfeeding among mothers attending government hospitals in Chennai. About 200 mothers who complied with the inclusion criteria were enrolled in this study. The participants were selected using the purposive sampling technique. A questionnaire for assessing knowledge, attitude, and practice of mothers on breastfeeding was prepared based on WHO and CDC guidelines. Iowa infant feeding attitude scale was used to assess the attitude variable questionnaire.

The questionnaire consisted of four parts which are as follows. Part I consisted of 20 questions with demographic details, obstetric details, and previous knowledge of breastfeeding. Part II comprised of 13 questions regarding the breastfeeding practices such as early initiation of breastfeeding, exclusive breastfeeding, use of pre and post-lacteal feeds, if colostrum was given, Bottle feeding practices, Rooming-in, and Skin to skin contact practices. Part III consisted of 15 questions, with questions on knowledge of breastfeeding, regarding colostrum, breastfeeding duration, mother's comfort, bonding between mother and newborn, neonatal and maternal health, maternal hygiene, and beauty. Questions were formulated in true or false format. The score was interpreted as 76 – 100% with good knowledge, 51 – 75% as fair knowledge, $\leq 50\%$ with poor knowledge.

For assessing the attitude towards breastfeeding, 17 questions were used. The responses were recorded as 0- Disagree, 1-Neutral, 2-Agree. These were interpreted as scores below 50% with a negative attitude towards breastfeeding and scored between 51 % - 100 % as a positive attitude towards breastfeeding. After obtaining Institutional ethical clearance, this study was conducted in a major government hospital in Chennai, Tamilnadu. Two hundred mothers of Infants attending paediatrics outpatient and inpatient were randomly selected. Data entry and data analysis were done with SPSS

software. All the data were tabulated and analysed. Demographic and obstetrical variables of mothers

were analysed in terms of frequency and percentage distribution.

Results

Table-1: Demographic characteristics of the mothers

Demographic details of a mother			
Demographic variable	Group	Frequency	Percent
Age in years	≤ 19	4	2
	20-25	84	42
	26-30	84	42
	≥ 31	28	14
Education	Illiterate	79	39.5
	Schooling	73	36.5
	Undergraduates	35	17.5
	Postgraduates	10	5
	Professional degree	3	1.5
Employment status	Government	5	2.5
	Private	35	17.5
	Business	4	2
	Homemaker	139	69.5
	Daily wages/coolie	17	8.5
Income in rupees	Below 5000	100	50
	5001-10000	57	28.5
	10001-15000	30	15
	15001-20000	10	5
	Above 20000	3	1.5
Religion	Hindu	141	70.5
	Christian	41	20.5
	Muslim	18	9
Family type	Nuclear family	131	65.5
	Joint family	69	34.5
Place of living	Urban	112	51.5
	Rural	97	48.5

Table 1 represents the frequency and percentage distribution of demographic variables of mothers and their infants. Concerning the maternal age, 4 (2%) were below 19 years and 84(42%) belonged to 20-25years. 84(42%) were in 26-30 years and 28(14%) were more than 31 years. Considering the educational status of mothers, 79 (39.5%) have completed up to middle school. 73(36.5%) have completed their high school. 35 (17.5%) have completed the undergraduate and 10 (5%) were post-graduates. Regarding the occupation of the mothers 5 (2.5%) of them were government employees, 35 (17.5%) were private employees, 4 (2%) were doing business, 139 (69.5%) were housewives and 17(8.5%) were daily wages/coolie. The family income of 100 (50%)


mothers was below Rs.5000, 57 (28.5%) was within the income group of Rs. 5001-10000, 30 (15%) belonged to the income group of Rs 10001-15000, 10 (5 %) was within the income group of Rs. 15001-20000, and 3 (1.5%) belonged to the income group of Rs above Rs.20000. Among the participants 141(70.5%) were Hindu, 41(20.5%) were Christian and 18(9.0%) belonged to the Muslim community. In regards to family type, 131 (65.5%) were living in a nuclear family and 69(34.5%) were living in a joint family.103(51.5 %) were from the urban area, 97(48.5%) were from a rural area. Among the 200 mothers, 105(52.5%) had one child,76(38%)had two children,19(9.5%) were multiparous.

Table-2: Children’s history

Details of last delivery of participated mother			
Age of the child	< 6month	138	69
	> 6month	62	31
Gender	Male	143	71.5
	Female	57	28.5
Type of delivery	Natural	105	52.5
	Less	95	47.5
Birth order	Single birth	195	97.5
	Twins	5	2.5
Gestational age at birth	Below 35 weeks	29	14.5
	35 to 38 weeks	171	85.5
Birth weight	≤ 2000 g	17	8.5
	2001 - 2500 g	44	22.0
	2501 - 3000 g	77	38.5
	≥ 3001g	62	31.0


From the above table, we can observe that 138(69%) of infants belonged less than 6 months, 62(31%) were more than 6 months of age. Among the children 143(71.5%) were male and the remaining were female babies. 105 (52.5%) mothers have delivered vaginally and 95 (47.5%) have undergone c-section. 171 (85.5%) mothers delivered at full-term, 29 (14.5%) was preterm delivery. 195(97.5%) were singleton birth, 5(2.5%) was twins. Around 17(8.5%) were LBW (low birth weight) with 44(22%) weighing 2001 -2500 grams, 77(38.5%) weighing 2501 -3000 grams and 62(31%) above 3001 grams.

Fig-1: Sources of information about breastfeeding



The greater number of participants obtained knowledge regarding breastfeeding from the nurses (59%), followed by physicians (29%). Around 12 % obtained by other sources.

Fig-2: Breastfeeding practices



The above figure showed that around 145 (72.5%) mothers have initiated breastfeeding within half an hour of delivery. Among the new-borns 14 (7%) of babies required postnatal admission. 7% of the neonates have received pre-lacteal feeds in the form of sugar water and honey. Only 181 (90.5%) of the babies have received the colostrum as their first feed. Around 19 (9.5%) were not fed with colostrum due to maternal hospitalization in other hospitals, the baby

was kept nil per oral due to some medical conditions. The use of prelacteal feeds along with colostrum was found from this study. Bottle-feeding practices accounted for 78(39.0%). 151(75.5%) babies were roomed in with the mothers within 30 minutes of delivery, others 49(24.5%) didn't have skin-to-skin contact with the mother due to delayed post-op recovery, and the baby got admitted to NICU. Rooming-in was practiced in 160(80%) of the babies

within an hour of birth while in 40(20%) rooming-in was not practiced as mother and child were separated due to admission of the neonate in NICU with some medical conditions like preterm birth, low birth

weight, asphyxia, hyperbilirubinemia and maternal complications such as delayed post-op recovery, maternal admission.

Table-3: KNOWLEDGE OF MOTHERS REGARDING BREASTFEEDING

KNOWLEDGE VARIABLE	TRUE		FALSE		MEAN	SD
	F	%	F	%		
Colostrum is the first breast milk	193	96.5	7	3.5	1.04	0.184241
Colostrum is important for the baby to maintain immunity	194	97	6	3	1.030	0.171015
Burping should be done after each feed	185	92.5	15	7.5	1.075	0.264052
Breastfeeding should be continued up to two years	144	72	56	28	1.28	0.450126
Exclusive breast milk can be given during the first six months	146	71	58	29	1.29	0.454901
Lactating mother should take healthy food to improve secretion of milk	168	84	32	16	1.16	0.367526
During breastfeeding, the mother should sit comfortably	146	73	54	27	1.27	0.445074
During breastfeeding, the mother should maintain eye to eye contact and talk with the baby	117	58.5	83	41.5	1.415	0.493958
Wash each breast with warm water before breastfeeding	141	70.5	59	29.5	1.415	0.493958
Awakening the baby while breastfeeding	121	60.5	79	39.5	1.295	0.457187
Breastfeeding helps in mother and child bonding	150	75	50	25	1.395	0.490077
Breastfeeding prevents diseases affecting breast	91	45.5	109	54.5	1.25	0.434099
Breastfeeding affects the beauty of feeding mothers	28	14	172	86	1.545	0.49922
The mother would not feed the child when she has diarrhoea	90	45	110	55	1.86	0.347858
Stop breastfeeding when you started weaning	71	35.5	129	64.5	1.645	0.479714

Knowledge of mothers regarding breastfeeding

In this study, 96.5% of mothers knew Colostrum being the initial milk and 194 (97%) mothers said Colostrum is necessary for the baby’s immunity. Regarding breastfeeding practices 185 (92.5%) mothers consented to burp after each feed is necessary. Among the participants, 144 (72%) reported breastfeeding to be practiced till two years and 142 (71%) reported EBF for the initial six months. Around 168 (84%) of mothers replied that lactating mothers should take healthy food to improve milk secretion and 146 (73%) of mothers

replied that during breastfeeding the mother should sit comfortably. 117(58.5%) of the participants stated that eye contact to be maintained during breastfeeding. Most consented that breastfeeding helps in maternal and child bonding. Almost 64.5% of mothers were against the statement of stopping breastfeeding when they start weaning. 91(45.5%) mothers knew as to breastfeeding reducing the risk of genital cancer. Among the participants, 70.5% of mothers had said to wash each breast with warm water before breastfeeding.

Table-4: Attitude of mothers towards breastfeeding

ATTITUDE SCORE	DISAGREE		NEUTRAL		AGREE		MEAN	SD
	F	%	F	%	F	%		
1The benefits of breast milk last only as long as the baby is breast fed	173	86.5	3	1.5	24	12.0	1.7450	.65737
2Formula Feeding Is More Convenient Than Breastfeeding	168	84.0	2	1.0	30	15.0	.3100	.71867
3breast feeding Increases Mother Infant Bonding	44	22.0	10	5.0	146	73.0	1.5100	.83269
4Breast milk is lacking in iron	104	52.0	16	8.0	80	40.0	1.1200	.95402
5FormuaFed Babies Are More Likely To Be Overfed Than Breastfed Babies	78	39.0	13	6.5	109	54.5	1.1550	.95685
6Formulafeeding is the better choice for the mother plans to go back to work	76	38.0	11	5.5	113	56.5	.8150	.95674
7Mother Who Formula Feed Miss One Of The Great Joys Of Motherhood	37	18.5	4	2.0	159	79.5	1.6100	.78164
8Women Should Not Breastfeed In Public Places Such As Restaurants	73	36.5	6	3.0	121	60.5	1.2400	.95759
9Breastfed Babies Are Healthier Than Fomula Fed Babies	31	15.5	6	3.0	163	81.5	1.6600	.73286
10Breastfed Babies Are More Likely To Be Overfed Than Formula Fed Babies	95	47.5	12	6.0	93	46.5	.9900	.97192
11Father feels left out if a mother breast feeds	72	36.0	19	9.5	109	54.5	.8150	.93549
12BreastMilk Is The Ideal Food For Babies	26	13.0	8	4.0	166	83.0	1.7000	.68729
13BreastMilk Is More Easily Digested Than Formula	33	16.5	9	4.5	158	79.0	1.6250	.75313
14 Formula is as healthy as breast milk	87	43.5	9	4.5	104	52.0	.9150	.97598
15Breast feeding Is More Convenient Than Fomula	36	18.0	9	4.5	155	77.5	1.5950	.77717
16Breast Milk Is Cheaper Than Formula	45	22.5	8	4.0	147	73.5	1.5100	.83870
17Mother who drinks occasionally should not breast feed	96	48.0	8	4.0	96	48.0	1.0000	.98225
	MEAN ATTITUDE SCORE						21.32	4.57168

Among the women 173 (86.5%) disagreed that the benefits of breastfeeding lasted only as long as the infant is breastfed. Nearly three-fourths of the mothers consented that breastfeeding is convenient rather than formula feeding. While 104 (52%) disagreed, 80 (40%) of them agreed to the statement “breast milk is lacking in iron”. Around 113 (56.5%) of the participants stated that “formula feeding is the better choice if the mother plans to go back to work”. Among the participants, 159 (79.5%) agreed that women who formula-fed miss the greatest joys of motherhood, and 109 (54.5%) of them agreed that

fathers feel left out if a mother breastfeeds. 60.5% agreed women can breastfeed in public places such as restaurants. Almost all of them stated that breast milk is ideal for babies 166 (83%), easily digested 158 (79%), and healthy for an infant than formula 163 (81.5%). 147 (73.5%) agreed that breast milk is affordable than Formula and 104 (52%) agreed that formula is as healthy as breast milk. Almost 96 (48%) felt that the mother who occasionally drinks alcohol should not breastfeed the baby. Overall the average score of the IIFAS (21.32 ± 4.57168, M±SD) lay in the range of ‘Negative breastfeeding attitudes’.

Table-5: Knowledge score

Knowledge score		
	Frequency	%
Poor (below 50%)	7	3.5
Fair (51% - 75%)	104	52
Good (76% - 100%)	89	44.5
Total	200	100

Most of the participants (52%) had fair knowledge, 45% of them had good knowledge, and very minimal(3%) had poor knowledge.

Table-6:Attitude score

Attitude score	Frequency	Percentage
Negative attitude (17 – 27)	3	1.5
A neutral attitude (28 – 39)	103	51.5
Positive attitude (40 ≤)	94	47.0

From this study, it was found that 51% of the mothers had a neutral attitude and 47% were having a positive attitude.

Table-7: Behaviour Score

Practice score	Frequency	Percentage
Non-desirable (below 50%)	52	26.0
Desirable (above 50%)	148	74.0
Total	200	100.0

Nearly three fourth of the participants (74%) had desirable practice scores.

Discussion

The first important step in breastfeeding is awareness and practices involved in breastfeeding. This study provides an insight into knowledge, attitude, and practices of breastfeeding in mothers hailing from an urban area, which could assist in intervention design and as a basis for more broad-based research in the other regions of the country. The Baby-Friendly Hospital Initiative (BFHI) was developed to encourage early initiation (i.e. within 30 minutes of delivery) of breastfeeding. Almost 72.5% of mothers initiated breastfed soon following birth in this study. This is comparatively greater than the study conducted by vijayalakshmi et al.,⁽⁶⁾ in Karnataka, India. The differences could be attributed to breastfeeding neonates kept nil per oral, preterm

birth, cleft lip and cleft palate, birth asphyxia, maternal exhaustion, delayed post-op recovery after LSCS, mothers hospitalized are some of the reasons for delayed breastfeeding initiation. Approximately 27 percent of them initiated breastfeeding after 30 minutes and the explanation for late initiation was due to LSCS and neonatal morbidity such as birth asphyxia. To ensure optimal growth and development, proper nutrition is deemed necessary during infancy. Breastfeeding contributes to short-term and long-term benefits, including protecting children from various acute and chronic disorders. The specific nutritional and antibody properties and benefits of colostrum are now established⁽⁷⁾. From the study, 90.5% of mothers fed colostrum, which is higher in comparison with other studies conducted in India⁽⁸⁾.EBF particularly in countries like India, is a

healthy, simple, economical, and emotionally satisfying means of feeding babies. Current research reported that approximately 93% have followed exclusive breastfeeding while the remaining 7% have historically given are-lacteal feed as sugar water, honey, donkey milk, etc. EBF in this study was 93%, which is greater than available national data (72%), while prelacteal feed was 7%, comparatively lower than the national data (28%)⁽²⁾. From this study, 39% of mothers practiced bottle feeding. Similarly, Adil Ali Nassir Ayed said that because of work, 56.7 percent of mothers adopted bottle feeding⁽⁹⁾. The justification of bottle-feeding practices was due to inadequate breast milk, working nature, alternative feeding. 44.5 percentage of mothers had strong awareness regarding breastfeeding. Around 97% of the participants knew about colostrum and its importance. In comparison, many studies found that 63.3% were informed of colostrum, and a study by Abroo Bashir et al showed almost identical results that 80% were informed of colostrum⁽¹⁰⁾.

Among the mothers, 92.5% practiced infant burping, which is greater than the other studies which reported 70% being aware of infant burping. About 72% thought they could continue breastfeeding till the second year. This finding is almost similar to a study by Vijayalakshmi et al⁽⁶⁾, which found that 68% knew they should continue breastfeeding till two years of age. Currently, the guidelines by WHO and UNICEF's Global Policy for Infant and Young Child Feeding states that children should be breastfed exclusively during the initial six months. Compared to Vijayalakshmi et al study, in the present survey, 85.2% knew EBF, which is comparatively poor. This result suggests raising awareness towards exclusive breastfeeding. ⁽⁶⁾Among the study participants 45.5% of mothers understood that breastfeeding can prevent breast-affecting disease, which is more than the previous survey by Abroo Bashir et al, which suggested that 17.6% of mothers recognized breastfeeding preventing breast-affecting disease⁽¹⁰⁾. Abroo Bashir et al⁽¹⁰⁾ also suggested 75% were aware that breastfeeding improves mother-child bonding, which is low compared to 87.9%. Most of the mothers were aware of breastfeeding practices. The mean attitude score in this study is 21.32 SD \pm 4.57, which points towards a negative attitude towards breastfeeding. The subjects of this study had better knowledge about breastfeeding but the attitude was

sub-optimal. This indicates that health educators should insist on breastfeeding and allow them to regularly apply their expertise to breastfeeding. Maternal variables and the medical condition of mothers causing a delayed initiation of feeding were post-op, maternal exhaustion, delayed post-op recovery after LSCS, admitted to other hospitals, delay in suturing episiotomy, and poor maternal feeding. New-born variables such as warmer infants, phototherapy, incubated, sustained under intensive care/observation unit and medical conditions such as neonates sustained in NPO due to preterm birth, heart issue, cleft lip and cleft palate, birth asphyxia were confirmed by previous studies⁽⁹⁾.

Conclusion

This study highlights the knowledge towards breastfeeding being fair among the urban population, but breastfeeding practices and attitudes were still sub-optimal. Few factors which were found from this study that could predict EBF includes the feeding experience and the employment of mothers. Insufficient breast milk was a major concern. Therefore, during their prenatal follow-up visits, healthcare providers should provide breastfeeding education to all women, especially women with low educational qualifications and no prior breastfeeding experience. Working towards resolving the barriers pertaining to breastfeeding, particularly for working mothers, alternatives must be provided including longer maternity leave and paid breaks for continued breastfeeding

References

1. World Health Organization. Global strategy for infant and young child feeding. World Health Organization; 2003.
2. Victora CG, Smith PG, Vaughan JP, Nobre LC, Lombardi C, Teixeira AM, Fuchs SC, Moreira LB, Gigante LP, Barros FC: Infant feeding and deaths due to diarrhea. A case-control study. *Am J Epidemiol.* 1989, 129: 1032-1041.
3. Arnold F, Parasuraman S, Arokiasamy P, Kothari M. Nutrition in India. National Family Health Survey (NFHS-3), India. 2005;6:59.
4. TurinCG Ochoa TJ The role of maternal breast milk in preventing infantile diarrhea in

- the developing world. *Curr Trop Med Rep* 2014;1:97-105
5. Martin, C.R.; Ling, P.-R.; Blackburn, G.L. Review of Infant Feeding: Key Features of Breast Milk and Infant Formula. *Nutrients* 2016, 8, 279
 6. Vijayalakshmi P, Susheela T, Mythili D. Knowledge, attitudes and breast-feeding practices of postnatal mothers: Across-sectionall survey. *Int J Health Sci.* 2015; 9(4):364-75
 7. Newburg DS. Innate immunity and human milk. *J Nutr* 2005;135(5):1308–12
 8. Pandey D, Sardana P, Saxena A, Dogra L, Coondoo A, Kamath A. Awareness and attitude towards breastfeeding among two generations of Indian women: a comparative study 2015. *PloS one.* 2015;10(5):1-9
 9. Ayed AAN. Knowledge, attitude, and practice regarding exclusive breastfeeding among mothers attending primary health care centers in Abha city. *Int J Med Sci Public Health* 2014;3:1355-1363.
 10. Bashir A, Mansoor S, Naikoo MY. Knowledge, attitude, and practices of postnatal mothers regarding breastfeeding: A cross-sectional study. *Int J Med Sci Public Health* 2018;7(9):725-730