



Cytomorphological Pattern Analysis Of Palpable And Usg – Guided Fnac Of Lymph Nodes – Study In A Tertiary Care Hospital

¹Dr(Maj) Manjunath HK, ²Dr Dharani VC, ³Dr Hema Rajkumari, ⁴Dr Vinitra K, ⁵Dr. Hassan Sona Rai, ⁶Dr Gudrun Koul, ⁷Dr Suma HV

¹Professor and HOD, ^{2,5}Associate Professor, ³2nd year Postgraduate, ^{4,6,7}Assistant Professor
Department of Pathology

***Corresponding Author:**

Dr. Hassan Sona Rai

BGS Medical College, Department of Pathology, No. 67, Uttarahalli main road, Kengeri, Bangalore, 560060, Karnataka, India

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract

Introduction :

The human body consists of approximately 600 lymph nodes. Tonsils, adenoids, spleen and Peyer's patches are various parts of the lymphoid tissue and their role is to provide immunity against various pathogens. Peripheral lymph nodes are located deep in the subcutaneous tissue and can be palpated if any thing causes them to enlarge. Lymph node enlargements are one of the commonest clinical presentation of patients and it encompasses a wide spectrum ranging from inflammation to a malignant lymphoma and finally metastatic malignancy. FNAC needle aspiration cytology (FNAC) is a rapid, simple, reliable, minimally invasive and cost-effective procedure which can be used as a outpatient setting however, histopathological examination remains as the gold standard in the evaluation of lymphadenopathy with ancillary tests..

Aim: The aim of the study was to evaluate the cytomorphological patterns of various diseases patterns in lymphadenopathy.

Materials And Methods :

A three years study of 100 cases of lymphadenopathy presenting to the Department of pathology from 1st January 2020 to 31st December 2022 taken up for our study. FNAC performed using a 23/24 gauge needle and 5ml syringe. Smears were fixed in 95% ethyl alcohol and stained with Hematoxylin and Eosin, Papanicolaou stain, Leishman stain and Giemsa stain. Special stains like periodic acid-Schiff for mucin (PAS) and Ziehl-Neelsen stain (ZN) stain for acid-fast bacilli (AFB), Reticulin stains, performed wherever required.

Results :

Most common lesion observed in our study was reactive lymphadenitis (25%), followed by granulomatous lymphadenitis (28.15%), tubercular lymphadenitis (17.39%), Neoplastic lesions(5%), metastatic lesions (14.23%). Cervical lymphadenopathy found to be the most common site in our study.

During the course of 3 years a total of 100 cases were received and were studied. Age of patients ranged from 1 year to 78 years. Females were 32 and males were 68 in number. Most common lesion found in our study was Reactive hyperplasia(25%), followed by Granulomatous (22%), Tuberculosis (20%), Suppurative (10%), Necrotic(5%), Parasitic (1%), Metastasis (12%), Lymphoma (5%).

Conclusion :

Our study highlighted the various cytomorphological patterns of lymphadenopathy. This study demonstrates that fine needle aspiration is a safe accurate and cost effective valuable screening tool in the evaluation of

various lymphadenopathies. FNAC analysis shows that accuracy is 95%, since histopathological diagnosis is invasive procedure.

Keywords: Lymph node, Lymphadenopathy, Fine needle aspiration cytology, Granulomatous, Malignant

Introduction

Lymphadenopathy is one of the most common clinical presentations of patients attending the outpatient department. The degree and pattern of morphological changes depend on the inciting stimulus and the intensity of the response. Thus, lymphadenopathy may be an incidental finding and / or primary or secondary manifestation of underlying diseases which may be neoplastic or non neoplastic.(1)Fine needle aspiration cytology(FNAC) as the first line of investigation has assumed importance in diagnosing a variety of disease processes as it is rapid, simple, reliable, minimally invasive and cost effective procedure which can be used in outpatient setting.(2) The etiology varies from an inflammatory process to a malignant condition. The knowledge of the pattern of lymphadenopathy in a given geographical region is essential for making a confident diagnosis or suspecting a disease. (3) Tuberculosis is the most common cause of lymphadenopathy in developing countries such as India and should be considered in every case of granulomatous lymphadenopathy unless proved otherwise. FNAC is a reliable, simple, safe, rapid and inexpensive method of establishing the diagnosis of lesions and masses at various sites and organs.(4)

FNAC has important role in the evaluation of peripheral lymphadenopathy, and it can be used as a safe alternative to excision biopsy. (5) FNAC has been used extensively for diagnosis of primary and secondary lymphadenopathy. The present study was undertaken to determine the role of FNAC in the evaluation of cytomorphological features of various lymph node lesions.

Materials And Methods :


This is a retrospective study composed of 100 cases of lymphadenopathy which was presented in the Department of Pathology for the past 3 years from 1st January 2020 to 31st December 2022, was taken up for this study. Informed consent was taken from all

patients in the study prior to the procedure. FNAC was performed under aseptic conditions, using a 22-24 gauge needle and 5 ml syringe. The smears thereafter were fixed in 95% ethyl alcohol and stained in various stains such as Hematoxylin and Eosin, Papanicolaou stain, Giemsa stain and Leishman stain. Cases of granulomatous disease, necrosis and suppuration, Ziehl- Neelsen stain was done for Gram negative acid fast bacilli. Few cases had fluid aspirate which was centrifuged and sediment smears was made using the above stains. The criteria for the lymph node aspirates to be diagnosed as tubercular lymphadenitis included the presence of epithelioid cell granuloma and caseous necrosis with or without Langhans giant cells or ZN positivity. Granulomatous lymphadenitis was diagnosed in the presence of epithelioid cell granuloma with or without giant cells and with the absence of necrosis.(6) This was followed by excision biopsy of the significantly enlarged lymph node from the same anatomical region was performed under local anaesthesia with the help of surgeons and the excised sample was sent for histomorphological examination in the same department.

Results :

A total of 100 aspirates were obtained out of which 68% were males and 32% were females. The age of the patients ranged from 1 year to 78 years with mean age of 39 years.

Maximum incidence of cases were seen in the age range of 30 – 40 years. A slight male preponderance with a ratio of 1.17 : 1 was noted.



Lymph nodes of varying sizes were subjected to FNAC. The smallest lymph node measured 1 cm and largest lymph node measured 3.5cm. Most of the lymph nodes ranged in size between 1 and 2 cm.

The most common group of lymph nodes aspirated were in a decreasing order cervical (38), submandibular (22), supraclavicular (18), axillary (12), and inguinal (10).

Cytomorphological diagnosis were made in all the 100 cases. Of the 100 cases, majority was reactive hyperplasia (30%), followed by tuberculosis (22% cases), granulomatous (15 %) , Suppurative (10 %), Necrotic (5 %), Parasitic (1%), metastatic malignancy (12 %) and lymphoma (5 %).

The tubercular lymph nodes were matted, firm in consistency, mobile and non-tender on palpation. For the cases of tuberculosis ZN staining was done and 6 cases showed acid fast bacilli. The rest of the cases were ZN negative, however CB-NAAT study showed positive for *Mycobacterium tuberculosis*.

Other lymph nodes were discrete, fixed with the overlying skin and non- tender. Non- Hodgkin's Lymphoma cases lymph nodes showed multi-regional involvement and were tender. Hodgkin's lymphoma cases lymph nodes also showed multi regional involvement and were tender on palpation.

The most common anatomical site of involvement of lymph nodes by different diseases was the cervical lymph nodes followed by submandibular and supraclavicular lymph node.

Discussion :

Lymph node aspiration is of great value in diagnosing lymphadenitis, lymphomas and metastatic carcinoma. The value of FNAC also lies in the early

direction of appropriate investigations, other than making the diagnosis. However, limitations and drawbacks of the procedure are there.

In developing countries like India, tuberculosis, acute upper respiratory tract infections and suppurative lymphadenitis are some of the common causes of lymphadenopathy. It has been stated that any significant lymph node enlargement not subsiding or remaining static in size for more than 2 weeks after conventional antibiotics need to be thoroughly investigated.

In this study, reactive lymphadenitis was the most common lesion, but in other studies the most common lesion recorded was tuberculosis. The next common lesion in the study was tuberculosis which was followed by metastatic malignancies in accordance with the study conducted by Malakaret al.(7) In many clinical settings, it is very difficult to decide which patient is more likely to have a reactive or neoplastic lymphadenopathy. Here, knowledge about the pattern of lymphadenopathy is helpful to the clinician for solving the dilemma.

Figure 1 : Microphotograph of reactive lymphadenitis showing mixed population of lymphoid cells.(400x)





Figure 2 : Microphotograph of tubercular epithelioid cell granuloma in the background of caseous necrosis.(400x).



Lymph node lesions can be seen in patients ranging from very early to advanced age. In our study, the youngest patient with lymphadenopathy was 1 year and oldest was 78 years old with a mean age of 39 years. In our study, male preponderance was noted with a male to female ratio of 1.2:1.(9)(10)

Table 1 :Showing distribution of age distribution of various lymphadenopathy.

Age(year s)	Reactive Hyperplasia	Granulomatous	Tuberculosis	Suppurative	Neoplastic
<1	-	-	-	-	-
1-10	-	2	4	3	1
11 -20	2	6	6	2	1
21 -30	3	7	4	5	2
31-40	15	5	3	-	-
41-50	5	2	3	-	1
51-60	-	-	-	-	-
61-70	-	-	-	-	-
71-80	-	-	-	-	-
TOTAL	25	22	20	10	5

Cervical lymph nodes were the most common group of lymph nodes 78 cases (78%) involved which was similar to that observed by Hirachand et al., Khajuria et al., and Chandanwale et al.(11)

Out of the 100 cases studied, 83 were benign/infectious and 17 cases were malignant. The most common cytological diagnosis was reactive hyperplasia (25%) in the present study. The criteria used to diagnose reactive hyperplasia was established included high cell density, polymorphic patterns of cells and considerable number of tingible bodies macrophages.

Similar findings was observed by Lakley et al. (12)(13)

Figure 3 : Microphotograph of Hodgkin’s lymphoma showing RS cell-Cytology, high power view(400x)





Figure 4 :Microphotograph of Large cell lymphoma




However, A B Pandav et al. (14) and A K Kochhar et al. (15) found Tuberculous lymphadenitis as the most common cytological diagnosis. These variations maybe due to sample size, age of patients, place and

duration of the study, local, genetic and environmental factors.

In our study, 22 cases of tuberculous lymphadenitis were subjected for ZN staining to confirm gram negative acid fast bacilli. 22 cases were positive for gram negative acid fast bacilli. Ng et al. (16) reported positivity in 41.6% cases. India is the country with the highest burden of tuberculosis that mainly involves the lungs followed by cervical lymph nodes. The portal of entry of TB bacilli into cervical lymph nodes is usually tonsils, or adenoids. This could be the reason for the high number of TB lymphadenitis in the present study. (17)

Figure 5: Microphotograph of Ziehl- Neelsen Stain showing gram negative acid fast bacilli.(100x)



12 cases of lymph nodes in our study showed malignant features after comparing with Histomorphological study. This is in correlation with the studies by Sharma R.I er al. (18) and Bhavani et al (19) where metastatic deposits were seen in 10.4% cases and 9.5% respectively.

Table 2: Showing the percentage distribution of lymph node lesions in decreasing order.

LESION	PERCENTAGE(%)
Cervical	38%
Submandibular	22%
Supraclavicular	18%
Axillary	12%

Inguinal	10%
----------	-----

Most of the metastatic deposits were from squamous cell carcinoma arising commonly in the tongue, alveolus, buccal mucosa, palate and from lung followed by adenocarcinoma.

This high percentage of squamous cell carcinoma was probably because of very high number of people have a bad habit of tobacco chewing. (20)

Cytology of lymph nodes has become a window for diagnosis of many disease conditions. Optimal material and experience, when combined, make cytological diagnosis of equal significance as histopathology. (21)However, it is not a replacement for conventional surgical pathology but is complimentary to identify the pathology of the disease.

We had one case(1%) which presented with multiple swelling over the right inguinal region measuring 2 x 3cm, clinically diagnosed as great saphenous-femoral junction thrombophlebitis. On ultrasound examination showed right sapheno-femoral junction incompetence. FNAC was done for right inguinal swelling. Microscopy showed slender microfilarial worms along with eggs, sheets of neutrophils along with lymphocytes, many plasma cells, eosinophils and cyst macrophages.

Filariasis is a term for infection caused by any of the filarial worms; however, in daily practice, the term filariasis is indicative of lymphatic filariasis caused by Wucheria or Brugia species. The vector used for transmission was the Culex mosquito. Microfilaria exhibit nocturnal periodicity in the peripheral blood. (22)Fine needle aspiration cytology can be a sensitive, cost-effective, and valuable tool for the detection of helminthic etiology in unexplained lymph node masses. (22,23)

Figure 6: Microphotograph of microfilaria(H&E)-high power view(400x).



Conclusion :

Fine needle aspiration cytology of lymph nodes is an excellent first line investigation to determine the nature of lesion. Accuracy of FNAC is comparable with histopathology. It is quick, safe, cost effective , minimally invasive and reliable and is readily accepted by the patient. It is useful screening tool in diagnosing both non- neoplastic and neoplastic lesions. ZN stain done in cases with granulomas, necrosis or suppuration is highly valuable for routine diagnosing of tuberculosis. It is an economical and convenient alternative to open biopsy of lymph nodes. With the introduction of FNAC, all the peripheral lymph nodes that are easily accessible can be assessed through a needle to arrive at the diagnosis.

The most frequent cases of lymphadenopathy are reactive lymphadenitis, tubercular lymphadenitis and metastatic malignancies. FNAC alone can aid in establishing the diagnosis in large number of cases. In certain situations it can be enough for diagnosis in clinical setting to avoid surgical procedure like biopsy. FNAC used in conjunction with clinical findings, radiological and laboratory investigations can be cost effective method.

References :

1. Pandit AA, Candes FP, Khubchandani SR. Fine needle aspiration cytology of lymph nodes. *J Postgrad Med* 1987;33:1344-6.
2. Pavithra P, Geetha JP. Role of fine needle aspiration cytology in the evaluation of the spectrum of lymph node lesions. *Int J Pharm Bio Sci* 2014;5:377-84.
3. Khajuria R, Goswami KC, Singh K, Dubey VK. Pattern of lymphadenopathy on fine needle aspiration cytology in Jammu. *JK Sci* 2006;8:157-9.
4. Kochhar AK, Duggal G, Singh K, Kochhar SK. Spectrum of cytological findings in patients with lymphadenopathy in rural population of South Haryana, India – Experience in a tertiary care hospital. *Internet J Pathol*2012;13:8.
5. Ajmal I, Imrann A. Comparison of FNAC vs. excision biopsy for suspected tuberculous cervical lymphadenopathy. *Ann King Edward Med Coll* 2003;9:216-8.
6. Hirachand S, Lakhey M, Akhter J, Thapa B. Evaluation of fine needle aspiration cytology of lymph nodes in Kathmandu Medical College, Teaching hospital. *Kathmandu Univ Med J(KUMJ)* 2009;7:139-42.
7. Malakar D, Jajoo IL, Swarup K, Gupta OP, Jain AP, Poflee VW. A clinical evaluation of fine needle aspiration cytology in the diagnosis of lymphadenopathy. *Indian J Tuberc*1991;38:17-9.
8. Das DK, Gupta SK. Fine needle aspiration cytodiagnosis of Hodgkin's disease and its subtypes, subtyping by differential counts. *Acta Cytol*1990;34:337-41.
9. Dhingra V, Misra V, Misra R, Bhatia R, Singhal M. Fine needle aspiration cytology(FNAC) as a diagnostic tool in paediatric lymphadenopathy. *J Clin Diagn Res* 2010; 4:2452-7.
10. Patra AK, Nanda BK, Mohapatra BK, Panda AK. Diagnosis of lymphadenopathy by fine needle aspiration cytology. *Indian J PatholMicrobiol*1983;26:273-8.
11. Khajuria R, Goswami KC, Singh K, Dubrey VK. Pattern of lymphadenopathy on fine needle aspiration cytology in Jammu. *JK Sci* 2006;8:157-9.
12. Lakhey M, Bhatta CP, Mishra S: Diagnosis of cytology, Acid fast Staining and Mantoux Test : *J Nepal Med As-soc* 2009;48(175):230-33.
13. Mamatha K. ArakeriSU : Utility of fine needle aspiration cytology (FNAC) in evaluation of cervical lymphadenopathy: *Indian Journal of Pathology and Oncology*, April – June 2017;(2):218-20.
14. A.B Pandav, P.P Patil, D. N. Lanjewar: cervical lymphadenopathy – Diagnosis by FNAC, A study of 219 cases: *Asian J Med Res* 2012;1(3):79-83.
15. A. K. Kochhar, G Duggal, K. Singh, S. K. Kochhar: Spectrum of cytological findings in

- patients with lymphadenopathy in rural population of southern Haryana, India – Experience in a tertiary care hospital. The internet journal of Pathology 2012; 13(2): 7-11.
16. Ng WF, Kung RT. Clinical research pathology of tuberculous lymphadenitis. A fine needle aspiration approach. J Honk Kong Med Assoc 1990; 42: 18-21.
 17. Arul P, Masilamani S, Akshatha C. Diagnostic efficacy of fine needle aspiration cytology in the evaluation of cervical lymphadenopathy. J Sci Soc 2016; 43:117-21.
 18. Sharma R.I, Dharaiya C.M. Study of fine needle aspiration cytology of lymphadenopathy in tertiary care centre of Ahmedabad, Gujarat. Trop J Path Micro 2018;(3):258-264.
 19. Bhavani C, Neeraja M, Varalakshmi KP, Ramana Babu PV. Chaitanya B, Sravani P. Role of FNAC in the diagnosis of cervical lymphadenopathy. Int J Med Res Rev 2014;2:599- 603.
 20. Joshi U, Modi B, Yadav S. A study on prevalence of chewing tobacco and existing quitting patterns in urban population of Jamnagar, Gujarat. DOI: 10.4103/0970-0218.62560.
 21. Shakya G, Malla S, Shakya K, Shrestha R. A study of FNAC of cervical lymph nodes. J Nepal Health Res Counc. 2009;7(1)1-5.
 22. Pathak B, Maimoon. Incidental finding of microfilaria in lymph node cytology:A case report. Cureus. 2022;doi10.7759/cureus.31275.
 23. Kesavan T M A, Narayan A, Rose F. An uncommon presentation of filariasis: A case report. IP International Journal of Medicine Paediatrics and Oncology.