



Predictive Value of Global Registry of Acute Coronary Events (GRACE) Score on acute heart failure : A hospital based cross-sectional study in Himachal Pradesh

Amit Kumar¹, Paramjot Kaur¹, R. C. Negi², Prem Machhan³, Jatinder Mokta³, Neeraj Ganju⁴

¹Junior Resident, ²Associate Professor, ^{3,4}Professor,
^{1,2,3}Department of Medicine, ⁴Department of Cardiology
Indira Gandhi Medical College, Shimla (H.P.)

***Corresponding Author:**

Amit Kumar

Junior Resident, Department of Medicine, Indira Gandhi Medical College, Shimla (H.P.)

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract

Introduction: Risk assessment and stratification plays an important role in management of Acute coronary syndrome (ACS). GRACE score is one of the most frequently used risk assessment tools. ACS can lead to heart failure (HF) during index hospitalization and we aimed to assess the performance of GRACE score in predicting in-hospital heart failure after index episode of MI.

Methods: We conducted a observational cross sectional study in a tertiary care hospital, analysing data from patients presenting with AMI. Demographic information, comorbidities, symptomatology were collected, GRACE score was calculated and correlated with left ventricular ejection fraction (LVEF) to assess their predictive value for acute HF during index MI presentation.

Results. A total of 283 patients were included in the study, with a mean age of 60.9 ± 12.0 years. Male patients constituted the majority, accounting for 74.9% of the total. 70.9% of patients with Grace score of ≤ 108 had preserved EF, whereas 55.1% of patients with score between 109 to 140 had preserved EF and the proportion of same for patients with GRACE score of more than 140 was only 33.8%. (p value- <0.001).

Conclusion: Our study emphasized the importance of comprehensive risk assessment and prognostic scoring using GRACE scoring model in predicting acute HF development. The GRACE score demonstrated a noteworthy correlation with LVEF, with increasing scores associated with a decreased proportion of patients with preserved EF.

Keywords: Acute myocardial infarction, Comorbidities, Heart failure, Left ventricular ejection fraction, Prognostic scores, GRACE score

Introduction

Acute myocardial infarction (AMI) is a well-known risk factor for Heart failure and similarly heart failure can complicate MI at admission or during hospital stay and severity of heart failure is associated with increased mortality [1-4]. Heart failure (HF) after myocardial infarction is the major cause of late morbidity, mortality and healthcare cost. The timing of heart failure after MI is important clinically, mechanistically and for research [5-6].

Three key time periods have to be defined, HF at the index MI presentation, during the course of the first admission, and post discharge [7]. several overlapping mechanisms contribute to HF after MI. HF during the index MI occurs due to a combination of myocardial stunning, myocyte necrosis, and acute mitral regurgitation due to papillary muscle dysfunction. HF during the hospitalisation may also be compounded by fluid or contrast overload, renal dysfunction, or complications such as ventricular

septal defect or cardiac tamponade. Late HF reflects the consequences of cardiomyocyte death and scar formation occurring alongside ventricular remodelling [8].

Many risk stratification models are now used including TIMI risk assessment model, GRACE model, Killip score etc. which predict severity of AMI and can be used as a predictor of heart failure, these markers tend to predict acute heart failure in hospital setting [9,10]. GRACE was developed involving more than 40 000 patients presenting with ACS as a clinical risk prediction tool for estimating the cumulative 6-month risk of death and death or myocardial infarction to facilitate triage and management of patients with ACS. GRACE is not restricted to any ST segment alterations.[11] Medical treatment of heart failure is based on type of heart failure and severity of heart failure. With evolving knowledge of MI and heart failure, early identification and initiation of therapy is important. Very limited studies have been conducted in hilly state of Himachal Pradesh, to study the profile of acute heart failure after acute myocardial infarction. Understanding the interplay between comorbidities, symptoms, and prognostic scores in AMI is crucial for optimizing patient management and improving outcomes, particularly in the context of heart failure (HF) development. The present study was planned to better understand the profile and prognostic indicators of heart failure patients of acute myocardial infarction during hospital stay in a tertiary care centre in Shimla, Himachal Pradesh.

Methodology:

This cross-sectional observational study was conducted at a tertiary care centre in Shimla, Himachal Pradesh over a period of one year, from January 1st, 2022, to December 31st, 2022. The patients admitted to the medicine and cardiology departments of the institute were considered for the study. Inclusion criteria encompassed individuals aged 18 years and above presenting with AMI, defined according to the fourth universal definition [12]. Exclusion criteria comprised patients below 18 years, individuals with prior myocardial infarctions, known heart failure, chronic kidney disease and individuals unwilling to give consent for the study.

Patients were provided with a comprehensive overview of the study's aims and objectives, and

those expressing willingness to participate were requested to provide written informed consent. Subsequently, a pre-designed pre-tested validated questionnaire was employed to gather information from the participants, ensuring that no identifiable details were recorded. The study encompassed comprehensive clinical evaluation, beginning with a detailed history to identify symptoms indicative of acute myocardial infarction, such as chest pain, dyspnea, syncope, and palpitations. A meticulous physical examination, encompassing anthropometric measurements and a systemic assessment with particular emphasis on detecting signs of cardiac failure. Biochemical investigations included high sensitivity Troponin I, complete hemogram, renal and liver function tests, HbA1c, BNP, lipid profile, thyroid profile, and other necessary tests as warranted. Electrocardiograms were conducted using a Heidelco Medicores HE 300 machine, assessing parameters like rate, rhythm, conduction disturbances, ST-T changes, and evidence of chamber enlargement. Echocardiographic examinations were performed using a Philips iE53 machine, with patients positioned supine or in a left decubitus position for optimal assessment. Monitoring of patients in wards involved close observation for symptoms and signs of acute heart failure, such as dyspnea, bibasal crepts, increasing jugular venous pulse, and gallop heart sounds, with repeated investigations conducted as deemed necessary. Anonymity of the patients and confidentiality of their responses was strictly maintained.

The collected data was entered into an MS Excel master sheet and subsequently tabulated and analyzed using statistical software, including OpenEpi version 3.01 and Statistical Package for Social Sciences (SPSS) version 22. Categorical data are represented as numbers and percentages (%), while quantitative data are expressed as mean and standard deviation. Analysis of categorical variables was done using Pearson's chi-square test and Fisher exact. A p value of <0.05 has been considered as statistically significant.


Results:

A total of 283 patients were included in the study, with a mean age of 60.9 ± 12.0 years. The majority of participants fell within the age bracket of 55 to 64

years (102 patients, 36%), followed by 79 patients (27.9%) aged between 65 to 74 years, and 48 patients (17%) aged 45 to 54 years. Male patients constituted the majority, accounting for 74.9% of the total. The most prevalent co-morbidity among the study cohort was hypertension, observed in 93 patients (32.9%),

closely followed by diabetes affecting a similar proportion of participants (88 patients, 31.1%). Smoking emerged as the most common modifiable risk factor, with over half of the participants being smokers (163 subjects, 57.6%), while peripheral artery disease was present in 47 patients (16.6%).

Figure 1: Comparison of GRACE score with LVEF among study participants



On comparison of Grace score and LVEF, it was observed that the proportion of patients with preserved EF reduced with increasing Grace score, i.e., 70.9% of patients with Grace score of ≤ 108 had preserved EF, whereas 55.1% of patients with score between 109 to 140 had preserved EF and the proportion of same for patients with Grace score of more than 140 was only 33.8%. This change in EF with Grace score was also found to be statistically significant with a p value of less than 0.001 (figure 1).

Discussion:

Over the past few decades, with advances in the management of acute MI, there has been a decline of over 40% in all-cause mortality during the acute phase of MI. AMI is the most common cause of HF which can present as early as in index hospitalization. Keeping this in mind, the present study was undertaken in the department of medicine/ cardiology in a tertiary care hospital of Shimla, Himachal Pradesh.

The initial assessment of left ventricular ejection fraction (LVEF) among acute myocardial infarction (AMI) cases holds paramount importance in guiding

patient management and prognostication. Our study, along with previous research, emphasizes the variability in LVEF distribution among different types and locations of MI. Understanding LVEF at presentation enables risk stratification and tailored therapeutic interventions.

The GRACE score, a risk assessment tool, exhibited a noteworthy correlation with LVEF in the present study. As the Grace score increased, the proportion of patients with preserved EF decreased significantly.

The findings of this study highlighted the association between left ventricular ejection fraction (LVEF) and GRACE score, underscoring its utility in risk stratification and guiding therapeutic decisions. Hence, there is a need for early risk assessment, prompt diagnosis, and tailored management strategies in improving outcomes for patients with AMI and acute HF.

References:

1. Wojtyniak B, Gierlotka M, Opolski G, Rabczenko D, Ozierański K, Gąsior M, Chlebus K, Wierucki Ł, Rutkowski D, Dziełak D, Poloński L. Observed and relative survival and 5-year outcomes of patients discharged after acute myocardial infarction: the nationwide AMI-PL database. *Kardiologia Polska (Polish Heart Journal)*. 2020;78(10):990-8.
2. Pagliaro BR, Cannata F, Stefanini GG, Bolognese L. Myocardial ischemia and coronary disease in heart failure. *Heart Failure Reviews*. 2020 Jan;25(1):53-65.
3. Groenewegen A, Rutten FH, Mosterd A, Hoes AW. Epidemiology of heart failure. *European journal of heart failure*. 2020 Aug;22(8):1342-56.
4. Liang KV, Pike F, Argyropoulos C, Weissfeld L, Teuteberg J, Dew MA, et al. Heart failure severity scoring system and medical- and health-related quality-of-life outcomes: the HEMO study. *Am J Kidney Dis*. 2011 Jul;58(1):84-92.
5. Jenča D, Melenovský V, Stehlik J, Staněk V, Kettner J, Kautzner J, et al. Heart failure after myocardial infarction: incidence and predictors. *ESC Heart Fail*. 2021 Feb;8(1):222-237.
6. Kim HY, Kim KH, Lee N, Park H, Cho JY, Yoon HJ, et al. Timing of heart failure development and clinical outcomes in patients with acute myocardial infarction. *Front Cardiovasc Med*. 2023;10 :1193973.
7. Cahill TJ, Kharbanda RK. Heart failure after myocardial infarction in the era of primary percutaneous coronary intervention: mechanisms, incidence and identification of patients at risk. *World journal of cardiology*. 2017 May 26;9(5):407.
8. Bahit MC, Kochar A, Granger CB. Post-myocardial infarction heart failure. *JACC: Heart failure*. 2018 Mar;6(3):179-86
9. Gawinski L, Engelseth P, Kozłowski R. Application of Modern Clinical Risk Scores in the Global Assessment of Risks Related to the Diagnosis and Treatment of Acute Coronary Syndromes in Everyday Medical Practice. *Int J Environ Res Public Health*. 2021 Aug 28;18(17):9103.
10. D'Ascenzo F, Biondi-Zoccai G, Moretti C, Bollati M, Omedè P, Sciuto F, et al. TIMI, GRACE and alternative risk scores in Acute Coronary Syndromes: A meta-analysis of 40 derivation studies on 216,552 patients and of 42 validation studies on 31,625 patients. *Contemp Clin Trials*. 2012;33(3):507–14.
11. Fox KA, Dabbous OH, Goldberg RJ, Pieper KS, Eagle KA, Van de Werf F, Avezum Á, Goodman SG, Flather MD, Anderson FA, Granger CB. Prediction of risk of death and myocardial infarction in the six months after presentation with acute coronary syndrome: prospective multinational observational study (GRACE). *bmj*. 2006 Nov 23;333(7578):1091.
12. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al. Fourth universal definition of myocardial infarction (2018). *Circulation*. 2018;138(20).